

# Vaccines for the National Immunisation Schedule



A reference card for vaccinators and other health professionals

our best protection

<p><b>Pregnancy</b></p>	 <p>Tdap injection (<b>Boostrix</b><sup>®</sup>)</p>	<p>+</p> <p>Influenza 1 injection (given annually) <b>Brand varies.</b></p>	
<p><b>6 Weeks</b></p>	 <p>RV1 oral vaccine (<b>Rotarix</b><sup>®</sup>)</p>	<p>+</p>  <p>DTaP-IPV-Hep B/Hib injection (<b>Infanrix</b><sup>®</sup> hexa)</p>	<p>+</p>  <p>PCV13 injection (<b>Prevenar 13</b><sup>®</sup>)</p> <p><b>From 1 Dec 2022</b></p>
<p><b>3 Months</b></p>	 <p>RV1 oral vaccine (<b>Rotarix</b><sup>®</sup>)</p>	<p>+</p>  <p>DTaP-IPV-Hep B/Hib injection (<b>Infanrix</b><sup>®</sup> hexa)</p>	
<p><b>5 Months</b></p>	 <p>DTaP-IPV-Hep B/Hib injection (<b>Infanrix</b><sup>®</sup> hexa)</p>	<p>+</p>  <p>PCV13 injection (<b>Prevenar 13</b><sup>®</sup>)</p> <p><b>From 1 Dec 2022</b></p>	
<p><b>12 Months</b></p>	 <p>MMR injection (<b>Priorix</b><sup>®</sup>)</p>	<p>+</p>  <p>PCV13 injection (<b>Prevenar 13</b><sup>®</sup>)</p> <p><b>From 1 Dec 2022</b></p>	
<p><b>15 Months</b></p>	 <p>Hib injection (<b>Hiberix</b><sup>®</sup>)</p>	<p>+</p>  <p>MMR injection (<b>Priorix</b><sup>®</sup>)</p>	<p>+</p>  <p>Varicella injection (<b>Varivax</b><sup>®</sup>)</p>
<p><b>4 Years</b></p>	 <p>DTaP-IPV injection (<b>Infanrix</b><sup>®</sup> IPV)</p>		
<p><b>11+12 Years</b></p>	 <p>Tdap injection (<b>Boostrix</b><sup>®</sup>)</p>	<p>+</p>  <p>HPV injection (<b>Gardasil</b><sup>®</sup> 9) (2 doses, 6 months apart)</p>	
<p><b>45 Years</b></p>	 <p>Tdap injection (<b>Boostrix</b><sup>®</sup>)</p>	<p><b>65 Years</b></p> <p>+</p>  <p>Tdap injection (<b>Boostrix</b><sup>®</sup>)</p> <p>+</p> <p>Influenza 1 injection (given annually). <b>Brand varies.</b></p>	<p>+</p>  <p>Shingles injection (<b>Shingrix</b><sup>®</sup>)</p>

# The National Immunisation Schedule

Age	Disease to protect against	Vaccine
Pregnancy	Tetanus + diphtheria + whooping cough (pertussis)	<b>Boostrix®</b>
	Influenza	Brand varies.
6 Weeks	Rotavirus (first dose must be given before 15 weeks)	<b>Rotarix®</b> (oral)
	Diphtheria + tetanus + whooping cough (pertussis) + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	<b>Infanrix® hexa</b>
	Pneumococcal disease	<b>Prevenar 13®</b>
3 Months	Rotavirus (second dose must be given before 25 weeks)	<b>Rotarix®</b> (oral)
	Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	<b>Infanrix® hexa</b>
5 Months	Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib)	<b>Infanrix® hexa</b>
	Pneumococcal disease	<b>Prevenar 13®</b>
12 Months	Measles + mumps + rubella	<b>Priorix®</b>
	Pneumococcal disease	<b>Prevenar 13®</b>
15 Months	<i>Haemophilus influenzae</i> type b (Hib)	<b>Hiberix®</b>
	Measles + mumps + rubella	<b>Priorix®</b>
	Chickenpox (varicella)	<b>Varivax®</b>
4 Years	Diphtheria + tetanus + whooping cough + polio	<b>Infanrix® IPV</b>
11+12 Years	Tetanus + diphtheria + whooping cough	<b>Boostrix®</b>
	Human papillomavirus (HPV)	<b>Gardasil®9</b> (2 doses, 6 months apart)
45 Years	Tetanus + diphtheria + whooping cough	<b>Boostrix®</b>
65 Years	Tetanus + diphtheria + whooping cough	<b>Boostrix®</b>
	Influenza	Brand varies.
	Shingles	<b>Shingrix®</b>