

# The National Immunisation Schedule

From – 1 JULY 2024

Stages	Disease to protect against	Vaccine
Pregnancy	Tetanus + diphtheria + whooping cough (pertussis)	Boostrix®
	Influenza	Brand varies
6 weeks	Rotavirus	Rotarix® (oral) (first dose must be given before 15 weeks)
	Diphtheria + tetanus + whooping cough (pertussis) + polio + hepatitis B + Haemophilus influenzae type b (Hib)	Infanrix® hexa
	Pneumococcal disease	Prevenar 13®
3 months	Rotavirus	Rotarix® (oral) (second dose must be given before 25 weeks)
	Diphtheria + tetanus + whooping cough + polio + hepatitis B + Haemophilus influenzae type b (Hib)	Infanrix® hexa
	Meningococcal B	Bexsero® (can be given at 8 weeks)
5 months	Diphtheria + tetanus + whooping cough + polio + hepatitis B + Haemophilus influenzae type b (Hib)	Infanrix® hexa
	Pneumococcal disease	Prevenar 13®
	Meningococcal B	Bexsero® (can be given at 4 months)
12 months	Measles + mumps + rubella	Priorix®
	Pneumococcal disease	Prevenar 13®
	Meningococcal B	Bexsero®
15 months	Haemophilus influenzae type b (Hib)	Act-HIB®
	Measles + mumps + rubella	Priorix®
	Chickenpox (varicella)	Varilrix®
4 years	Diphtheria + tetanus + whooping cough + polio	Infanrix® IPV
9-13 years	Human papillomavirus (HPV)	Gardasil®9 (2 doses, 6 months apart)
	Tetanus + diphtheria + whooping cough	Boostrix® (from 11 years)
45 years	Tetanus + diphtheria + whooping cough	Boostrix®
65 years	Tetanus + diphtheria + whooping cough	Boostrix®
	Influenza	Brand varies
	Shingles	Shingrix® (2 doses, 2 - 6 months apart)

# Vaccines for the National Immunisation Schedule

A reference card for vaccinators and other health professionals

From – 1 JULY 2024

Pregnancy		Tdap injection <b>(Boostrix®)</b> Every pregnancy from 13 weeks		Influenza 1 injection (given annually) Brand varies		
6 weeks		+ RV1 oral vaccine (Rotarix®)		+ DTaP-IPV-Hep B/Hib injection (Infanrix® hexa)		+ PCV13 injection (Prevenar 13®)
3 months		+ RV1 oral vaccine (Rotarix®)		+ DTaP-IPV-Hep B/Hib injection (Infanrix® hexa)		+ MenB injection (Bexsero®) Can be given at 8 weeks
5 months		+ DTaP-IPV-Hep B/Hib injection (Infanrix® hexa)		+ PCV13 injection (Prevenar 13®)		+ MenB injection (Bexsero®) Can be given at 4 months
12 months		+ MMR injection (Priorix®)		+ PCV13 injection (Prevenar 13®)		+ MenB injection (Bexsero®)
15 months		+ Hib injection (Act-HIB®)		+ MMR injection (Priorix®)		+ Varicella injection (Varilrix®)
4 years		DTaP-IPV injection (Infanrix® IPV)				
9-13 years		HPV injection (Gardasil® 9) 2 doses, 6 months apart		+ Tdap injection (Boostrix®) From 11 years		
45 years		+ Tdap injection (Boostrix®)		+ Influenza 1 injection (given annually) Brand varies		+ Shingles injection (Shingrix®) 2 doses, 2 – 6 months apart
65 years		+ Tdap injection (Boostrix®)				